arXiv:1609.08453v1 [math.GM] 23 Sep 2016

Generalized Weyl Conformal Curvature Tensor of a Generalized Riemannian Space

Nenad O. Vesić*

Abstract

It is generalized Weyl conformal curvature tensor in the case of a conformal mappings of a generalized Riemannian space in this paper. Moreover, it is found universal generalizations of it without any additional assumption. A method used in this paper may help different scientists in their researching.

1 Introduction

A lot of researchers have studied, developed and applied Weyl curvature tensor and Weyl conformal curvature tensor respectively defined as (see [1–3, 5, 7, 11, 13]):

$$\begin{aligned} C^i_{jmn} = R^i_{jmn} &+ \frac{1}{N-2}\big(\delta^i_n R_{jm} - \delta^i_m R_{jn} + g_{jm}R^i_n - g_{jn}R^i_m\big) \\ &+ \frac{1}{(N-1)(N-2)}\big(\delta^i_m g_{jn} - \delta^i_n g_{jm}\big)R, \end{aligned} \tag{1.1}$$

$$\begin{aligned} C_{ijmn} = R_{ijmn} &+ \frac{1}{N-2}\big(g_{in}R_{jm} - g_{im}R_{jn} + g_{jm}R_{in} - g_{jn}R_{im}\big) \\ &+ \frac{1}{(N-1)(N-2)}\big(g_{im}g_{jn} - g_{in}g_{mj}\big)R, \end{aligned} \tag{1.2}$$

of a Riemannian space $\mathbb{R}_N$endowed with a metric tensor g_{ij} symmetric by indices i and j and $R^i_j = g^{i\alpha}R_{\alpha j}$ in their researches. Some of them who

*Faculty of Science and Mathematics Niš

Key words: Weyl conformal curvature tensor, conformal mapping, Riemannian and generalized Riemannian space, invariant

2010 **Math. Subj. Classification:** 14L24, 14A22, 53A30, 53B20

were focused on this tensor of a Riemannian space $\mathbb{R}_N$ are H. M. Abood [1], S. Bochner [2], E.Goulart and M. Novello [5] and many others. A magnitude R_{ijmn} in this formula is a covariant curvature tensor of the space $\mathbb{R}_N$ defined as

$$R_{ijmn} = g_{i\alpha} R^{\alpha}_{jmn} \tag{1.3}$$

for a curvature tensor R^{i}_{jmn} of the space $\mathbb{R}_N$, for the corresponding Ricci tensot $R_{ij} = R^{\alpha}_{ij\alpha}$ and for the corresponding scalar curvature $R = g^{\alpha\beta} R_{\alpha\beta}$.

1.1 Some applications of a Weyl conformal curvature tensor

Goulart and Novello (see [5]) applied a Weyl conformal curvature tensor (**??**) in their research about a cosmological stability of it. In mathematics, S. Bochner [2] applied a Weyl conformal curvature tensor for his researches about N-dimensional spaces (even dimensional ones, the case $N = 2k$) endowed with a Hermitian metrics

$$ds^2 = 2 g_{\alpha\beta^*} dz_\alpha d\overline{z}_\beta$$

which satisfy a Kähler assumption

$$\frac{\partial g_{\alpha\gamma^*}}{\partial z_\beta} = \frac{\partial g_{\beta\gamma^*}}{\partial z_\alpha}$$

and he causally defined a tensor

$$\begin{aligned} B_{ij^*mn^*} &= R_{ij^*mn^*} \\ &\quad - \frac{1}{k+2}\big(g_{ij^*} R_{mn^*} + g_{in^*} R_{mj^*} + g_{mj^*} R_{in^*} + g_{mn^*} R_{ij^*}\big) \\ &\quad + \frac{R}{2(k+1)(k+2)}\big(g_{ij^*} g_{mn^*} + g_{in^*} g_{j^*m}\big). \end{aligned} \tag{1.4}$$

1.2 Motivation

The German mathematical physicist Hermann Weyl (1885-1955) made many important contributions to physics. It is well known a Weyl projective tensor (an important magnitude for a force of gravity researching). H. Weyl wanted to know does it exist a tensor similar to the curvature tensor of a Riemannian space by its symmetry properties but invariant under conformal mappings.

A tensor such that exists and it is named Weyl conformal curvature tensor. This tensor is given in the equation (**??**).

The primary aim of this paper is to generalize the Weyl conformal curvature tensor in the case of a generalized Riemannian space. Moreover, different authors restrict their research about non-symmetric affine connected spaces onto special cases. One of these special cases is equitorsion mappings, i.e. mappings which save the non-symmetric part $\Gamma^{i}_{\underset{\vee}{jk}}$ of affine connection coefficients (see [14]). The aim of this paper is to exclude all additional assumptions which have been considered as necessary ones for research about conformal mappings.

2 A brief recall of generalized Riemannian spaces

An N-dimensional manifold $\mathcal{M}_N$ endowed with a metric tensor $g_{ij} = g_{ij}(x)$ (a continually differentiable function at points of the manifold $\mathcal{M}_N$) non-symmetric by indices i and j is said to be **the generalized Riemannian space** $\mathbb{GR}_N$ [4, 8]. A symmetric and an anti-symmetric part of the metric tensor g_{ij} are respectively defined as

$$g_{\underline{ij}} = \frac{1}{2}\big(g_{ij} + g_{ji}\big) \quad \text{and} \quad g_{\underset{\vee}{ij}} = \frac{1}{2}\big(g_{ij} - g_{ji}\big). \tag{2.1}$$

Remark 2.1 *An anti-symmetrization without division is going to be denoted with square brackets as*

$$X_{\ldots[i\ldots j]\ldots} = X_{\ldots i\ldots j\ldots} - X_{\ldots j\ldots i\ldots},$$

for an indexed magnitude X.

The corresponding determinant

$$g = \begin{vmatrix} g_{\underline{11}} & \cdots & g_{\underline{1N}} \\ \vdots & \ddots & \vdots \\ g_{\underline{N1}} & \cdots & g_{\underline{NN}} \end{vmatrix} \tag{2.2}$$

is a metric determinant of the space $\mathbb{GR}_N$. Specially interesting for researchers is the case of $g \neq 0$ (**regular generalized Riemannian space**).

Remark 2.2 *It is going to be studied generalized Riemannian spaces with a metric determinant $g > 0$. A case $g < 0$ may be obtained with a sign-change $\tilde{g}_{i1} = -g_{i1}$;*

Magnitudes

$$\Gamma_{i.jk} = \frac{1}{2}\big(g_{ji,k} - g_{jk,i} + g_{ik,j}\big) \quad \text{and} \quad \Gamma^i_{jk} = g^{\underline{\alpha i}}\Gamma_{\alpha.jk} \tag{2.3}$$

are a first and a second kind Christoffel symbol of the space $\mathbb{GR}_N$. Second kind Christoffel symbols are affine connection coefficients of the space $\mathbb{GR}_N$.

A space $\mathbb{R}_N$ with affine connection coefficients $\Gamma^i_{\underline{jk}} = \frac{1}{2}\big(\Gamma^i_{jk} + \Gamma^i_{kj}\big)$ is said to be the associated space of the space $\mathbb{GR}_N$.

There is just one type of a covariant differentiation based on an affine connection of a Riemannian space $\mathbb{R}_N$. This type covariant derivative of a tensor a^i_j by x^k is defined as

$$a^i_{j;k} = a^i_{j,k} + \Gamma^i_{\underline{\alpha k}}a^\alpha_j - \Gamma^\alpha_{\underline{jk}}a^i_\alpha, \tag{2.4}$$

for a partial derivative denoted with comma.

For the reason of only one Ricci-type identity in this case, it exists just one curvature tensor of the associated space $\mathbb{R}_N$ defined as

$$R^i_{jmn} = \Gamma^i_{\underline{jm};n} - \Gamma^i_{jn;m} + \Gamma^\alpha_{\underline{jn}}\Gamma^i_{\underline{\alpha m}} - \Gamma^\alpha_{\underline{jm}}\Gamma^i_{\underline{\alpha n}}. \tag{2.5}$$

It is proved (see [6]) an anti-symmetric part $\underset{\vee}{g}_{ij}$ of a metric tensor of a generalized Riemannian space $\mathbb{GR}_N$ may be expressed in the form

$$\underset{\vee}{g}_{ij} = A^\alpha_i g_{\underline{\alpha j}}, \tag{2.6}$$

for a tensor A^i_j of the type (1, 1). From this equation it directly holds (the quotient rule) the tensor A^i_j has the form

$$A^i_j = g^{\underline{i\alpha}}\underset{\vee}{g}_{j\alpha}. \tag{2.7}$$

Before that, it had been involved (see [9]) four types of covariant derivation with regard of a non-symmetric affine connection. Caused by that it was obtained (see [8]) there is more than one curvature tensor of a non-symmetric affine connection space.

There are five linearly independent curvature tensors of a non-symmetric affine connection space [10]. In the case of a generalized Riemannian space $\mathbb{GR}_N$, these curvature tensors are:

$$\underset{1}{R}{}^i_{jmn} = R^i_{jmn} + \underset{\vee}{\Gamma}{}^i_{jm;n} - \underset{\vee}{\Gamma}{}^i_{jn;m} + \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}{}^i_{\alpha n} - \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}{}^i_{\alpha m}, \tag{2.8}$$

$$\underset{2}{R}{}^i_{jmn} = R^i_{jmn} - \underset{\vee}{\Gamma}{}^i_{jm;n} + \underset{\vee}{\Gamma}{}^i_{jn;m} + \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}{}^i_{\alpha n} - \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}{}^i_{\alpha m}, \tag{2.9}$$

$$\underset{3}{R}{}^i_{jmn} = R^i_{jmn} + \underset{\vee}{\Gamma}{}^i_{jm;n} + \underset{\vee}{\Gamma}{}^i_{jn;m} - \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}{}^i_{\alpha n} + \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}{}^i_{\alpha m} - 2\underset{\vee}{\Gamma}{}^\alpha_{mn}\underset{\vee}{\Gamma}{}^i_{\alpha j}, \tag{2.10}$$

$$\underset{4}{R}{}^i_{jmn} = R^i_{jmn} + \underset{\vee}{\Gamma}{}^i_{jm;n} + \underset{\vee}{\Gamma}{}^i_{jn;m} - \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}{}^i_{\alpha n} + \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}{}^i_{\alpha m} + 2\underset{\vee}{\Gamma}{}^\alpha_{mn}\underset{\vee}{\Gamma}{}^i_{\alpha j}, \tag{2.11}$$

$$\underset{5}{R}{}^i_{jmn} = R^i_{jmn} - \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}{}^i_{\alpha n} + \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}{}^i_{\alpha m}. \tag{2.12}$$

3 Generalized Weyl conformal curvature tensor

Let $\mathbb{GR}_N$ be a regular generalized Riemannian space. A Weyl conformal curvature tensor $\underline{C}_{ijmn}$ of a space $\mathbb{R}_N$ associated to the space $\mathbb{GR}_N$ has the form

$$\begin{aligned}\underline{C}_{ijmn} = R_{ijmn} &- \frac{1}{N-2}\big(g_{\underline{jm}}R_{in} - g_{\underline{jn}}R_{im} + g_{\underline{in}}R_{jm} - g_{\underline{im}}R_{jn}\big)\\ &- \frac{R}{(N-1)(N-2)}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big).\end{aligned} \tag{3.1}$$

From curvature tensors (**??**-**??**) one obtains covariant curvature tensors $\underset{k}{R}_{ijmn} = g_{\underline{i\alpha}}\underset{k}{R}{}^\alpha_{jmn}, k = 1, \ldots, 5$, are:

$$\underset{1}{R}_{ijmn} = R_{ijmn} + \underset{\vee}{\Gamma}_{i.jm;n} - \underset{\vee}{\Gamma}_{i.jn;m} + \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}_{i.\alpha n} - \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}_{i.\alpha m}, \tag{3.2}$$

$$\underset{2}{R}_{ijmn} = R_{ijmn} - \underset{\vee}{\Gamma}_{i.jm;n} + \underset{\vee}{\Gamma}_{i.jn;m} + \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}_{i.\alpha n} - \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}_{i.\alpha m}, \tag{3.3}$$

$$\begin{aligned}\underset{3}{R}_{ijmn} = R_{ijmn} &+ \underset{\vee}{\Gamma}_{i.jm;n} + \underset{\vee}{\Gamma}_{i.jn;m}\\ &- \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}_{i.\alpha n} + \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}_{i.\alpha m} - 2\underset{\vee}{\Gamma}{}^\alpha_{mn}\underset{\vee}{\Gamma}_{i.\alpha j},\end{aligned} \tag{3.4}$$

$$\begin{aligned}\underset{4}{R}_{ijmn} = R_{ijmn} &+ \underset{\vee}{\Gamma}_{i.jm;n} + \underset{\vee}{\Gamma}_{i.jn;m}\\ &- \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}_{i.\alpha n} + \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}_{i.\alpha m} + 2\underset{\vee}{\Gamma}{}^\alpha_{mn}\underset{\vee}{\Gamma}_{i.\alpha j},\end{aligned} \tag{3.5}$$

$$\underset{5}{R}_{ijmn} = R_{ijmn} - \underset{\vee}{\Gamma}{}^\alpha_{jm}\underset{\vee}{\Gamma}_{i.\alpha n} + \underset{\vee}{\Gamma}{}^\alpha_{jn}\underset{\vee}{\Gamma}_{i.\alpha m}. \tag{3.6}$$

A Weyl conformal curvature tensor $\underline{C}_{ijmn}$ is obtained to be an invariant of a conformal mapping

$$f : \mathbb{R}_N \to \overline{\mathbb{R}}_N$$

determined with an equation of the type

$$\overline{g}_{\underline{ij}} = e^{2\psi} g_{\underline{ij}} \tag{3.7}$$

for metric tensors $g_{\underline{ij}}$ and $\overline{g}_{\underline{ij}}$ of associated spaces $\mathbb{R}_N$ and $\overline{\mathbb{R}}_N$ of generalized Riemannian spaces $\mathbb{GR}_N$ and $\mathbb{G}\overline{\mathbb{R}}_N$ respectively.

It was generalized and studied the conformal mapping theory of a generalized Riemannian space $\mathbb{GR}_N$ (see [14]). These mappings are generated with an equation of the type

$$\overline{g}_{ij} = e^{2\psi} g_{ij}, \tag{3.8}$$

for a continually-differentiable function. A direct corollary of the equation (**??**) is

$$\overline{g}^{\underline{ij}} = e^{-2\psi} g^{\underline{ij}}. \tag{3.9}$$

From these results it directly holds next equalities are valid ones:

$$\overline{g}^{\underline{ij}} \overline{g}_{mn} = e^{-2\psi} g^{\underline{ij}} e^{2\psi} g_{mn} = g^{\underline{ij}} g_{mn}, \tag{3.10}$$

i.e. a tensor A^i_j from the equation (**??**) is an invariant of a conformal mapping f between generalized Riemannian spaces.

Using the fact $g^{\underline{ij}}{}_{;k} = 0$, the equation (**??**) and the expressions of a second kind Christoffel symbol, one obtains an anti-symmetric part $\Gamma^i_{\underset{\vee}{jk}} = \frac{1}{2}\big(\Gamma^i_{jk} - \Gamma^i_{kj}\big)$ of a second kind Christoffel symbol Γ^i_{jk} has a form

$$\Gamma^i_{\underset{\vee}{jk}} = \frac{1}{2}\big(A^i_{j;k} - A^i_{k;j} - g^{\underline{i\alpha}} g_{\underset{\vee}{jk};\alpha}\big). \tag{3.11}$$

From this equation one obtains a following equality holds:

$$\begin{aligned} \overline{\Gamma}^i_{\underset{\vee}{jk}} - \Gamma^i_{\underset{\vee}{jk}} = \frac{1}{2}\Big(\big(\overline{\Gamma}^i_{\underline{\alpha k}} - \Gamma^i_{\underline{\alpha k}}\big) A^\alpha_j - \big(\overline{\Gamma}^i_{\underline{\alpha j}} - \Gamma^i_{\underline{\alpha j}}\big) A^\alpha_k\Big) \\ - \frac{1}{2}\big(\overline{g}^{\underline{i\alpha}} \overline{g}_{\underset{\vee}{jk};\alpha} - g^{\underline{i\alpha}} g_{\underset{\vee}{jk};\alpha}\big), \end{aligned} \tag{3.12}$$

or equivalently

$$\underset{1}{\mathcal{I}}{}^i_{jk} = \underset{1}{\overline{\mathcal{I}}}{}^i_{jk},$$

for

$$\underset{1}{\mathcal{I}}{}^{i}_{jk} = \underset{\vee}{\Gamma}{}^{i}_{jk} + \frac{1}{2}\big(\Gamma^{i}_{\underline{\alpha j}} A^{\alpha}_{k} - \Gamma^{i}_{\underline{\alpha k}} A^{\alpha}_{j} + g^{\underline{i\alpha}} \underset{\vee}{g}_{jk;\alpha}\big). \tag{3.13}$$

First and second kind Christoffel symbols $\Gamma_{i.jk}$ and Γ^{i}_{jk} under a conformal mapping $f : \mathbb{GR}_N \to \mathbb{G}\overline{\mathbb{R}}_N$ satisfy rules:

$$\overline{\Gamma}_{i.jk} = e^{2\psi}\big(\Gamma_{i.jk} + g_{ji}\psi_k - g_{jk}\psi_i + g_{ik}\psi_j\big), \tag{3.14}$$

$$\overline{\Gamma}^{i}_{jk} = \Gamma^{i}_{jk} + \psi_j \delta^{i}_{k} + \psi_k \delta^{i}_{j} - \psi_\alpha g^{\underline{\alpha i}} g_{\underline{jk}} + \xi^{i}_{jk} \tag{3.15}$$

for $\psi_j = \partial\psi/\partial x^j = \psi_{,j}$ and $\xi^{i}_{jk} = \underset{\vee}{\overline{\Gamma}}{}^{i}_{jk} - \underset{\vee}{\Gamma}{}^{i}_{jk}$.

From the equation (**??**) it directly holds (a contraction $i = k$) the expression

$$\psi_j = \frac{1}{N}\big(\overline{\Gamma}^{\alpha}_{\underline{j\alpha}} - \Gamma^{\alpha}_{\underline{j\alpha}}\big) \tag{3.16}$$

is a valid one.

The equation (**??**) confirm the validity of a following equality:

$$\begin{aligned}\underset{\vee}{\overline{\Gamma}}{}_{i.jk} = e^{2\psi}\underset{\vee}{\Gamma}{}_{i.jk} - \frac{1}{N}\big(\underset{\vee}{\overline{g}}{}_{ij}\overline{\Gamma}^{\alpha}_{\underline{k\alpha}} - \underset{\vee}{\overline{g}}{}_{ik}\overline{\Gamma}^{\alpha}_{\underline{j\alpha}} + \underset{\vee}{\overline{g}}{}_{jk}\overline{\Gamma}^{\alpha}_{\underline{i\alpha}}\big) \\ + \frac{1}{N}e^{2\psi}\big(\underset{\vee}{g}{}_{ij}\Gamma^{\alpha}_{\underline{k\alpha}} - \underset{\vee}{g}{}_{ik}\Gamma^{\alpha}_{\underline{j\alpha}} + \underset{\vee}{g}{}_{jk}\Gamma^{\alpha}_{\underline{i\alpha}}\big).\end{aligned} \tag{3.17}$$

From the definition of a conformal mapping it directly holds determinants $g = \det\|g_{\underline{ij}}\|$ and $\overline{g} = \det\|\overline{g}_{\underline{ij}}\|$ satisfy a relation

$$\overline{g} = e^{2N\psi} g$$

which proves it holds an equation

$$\psi = \frac{1}{2N}\ln\frac{\overline{g}}{g}. \tag{3.18}$$

Involving this result into the equation (**??**) one proves a following equality holds:

$$\begin{aligned}\overline{\Gamma}_{\underset{\vee}{i.jk}} &= \Gamma_{\underset{\vee}{i.jk}} + \big(1-\overline{g}^{-\frac{1}{N}}\big)\overline{\Gamma}_{\underset{\vee}{i.jk}} - \big(1-g^{-\frac{1}{N}}\big)\Gamma_{\underset{\vee}{i.jk}} \\ &- \frac{1}{N\overline{g}^{\frac{1}{N}}}\big(\overline{g}_{ij}\overline{\Gamma}^{\alpha}_{\underline{k\alpha}} - \overline{g}_{\underset{\vee}{ik}}\overline{\Gamma}^{\alpha}_{\underline{j\alpha}} + \overline{g}_{\underset{\vee}{jk}}\overline{\Gamma}^{\alpha}_{\underline{i\alpha}}\big) \\ &+ \frac{1}{Ng^{\frac{1}{N}}}\big(g_{ij}\Gamma^{\alpha}_{\underline{k\alpha}} - g_{\underset{\vee}{ik}}\Gamma^{\alpha}_{\underline{j\alpha}} + g_{\underset{\vee}{jk}}\Gamma^{\alpha}_{\underline{i\alpha}}\big),\end{aligned} \tag{3.19}$$

or equivalently

$$\underset{2}{\mathcal{I}}_{ijk} = \underset{2}{\overline{\mathcal{I}}}_{ijk},$$

for

$$\underset{2}{\mathcal{I}}_{ijk} = \Gamma_{\underset{\vee}{i.jk}} - \big(1-g^{-\frac{1}{N}}\big)\Gamma_{\underset{\vee}{i.jk}} + \frac{1}{Ng^{\frac{1}{N}}}\big(g_{ij}\Gamma^{\alpha}_{\underline{k\alpha}} - g_{\underset{\vee}{ik}}\Gamma^{\alpha}_{\underline{j\alpha}} + g_{\underset{\vee}{jk}}\Gamma^{\alpha}_{\underline{i\alpha}}\big). \tag{3.20}$$

Let a conformal mapping $f : \mathbb{GR}_N \to \mathbb{G}\overline{\mathbb{R}}_N$ be determined with an equation of the type

$$\overline{\Gamma}^{i}_{jk} = \Gamma^{i}_{jk} + P^{i}_{jk}, \tag{3.21}$$

for $P^{i}_{jk} = \overline{\Gamma}^{i}_{jk} - \Gamma^{i}_{jk}$. In this case we have the following equalities are satisfied:

$$\begin{aligned}\underset{2}{\overline{\mathcal{I}}}_{ijm\bar{;}n} &= \underset{2}{\overline{\mathcal{I}}}_{ijm,n} - \overline{\Gamma}^{\alpha}_{\underline{in}}\underset{2}{\overline{\mathcal{I}}}_{\alpha jm} - \overline{\Gamma}^{\alpha}_{\underline{jn}}\underset{2}{\overline{\mathcal{I}}}_{i\alpha n} - \overline{\Gamma}^{\alpha}_{\underline{mn}}\underset{2}{\overline{\mathcal{I}}}_{ij\alpha} \\ &= \underset{2}{\mathcal{I}}_{ijm,n} - \big(\Gamma^{\alpha}_{\underline{in}} + P^{\alpha}_{\underline{in}}\big)\underset{2}{\mathcal{I}}_{\alpha jm} - \big(\Gamma^{\alpha}_{\underline{jn}} + P^{\alpha}_{\underline{jn}}\big)\underset{2}{\mathcal{I}}_{i\alpha n} - \big(\Gamma^{\alpha}_{\underline{mn}} + P^{\alpha}_{\underline{mn}}\big)\underset{2}{\mathcal{I}}_{ij\alpha} \\ &= \underset{2}{\mathcal{I}}_{ijm;n} - P^{\alpha}_{\underline{in}}\underset{2}{\mathcal{I}}_{\alpha jm} - P^{\alpha}_{\underline{jn}}\underset{2}{\mathcal{I}}_{i\alpha n} - P^{\alpha}_{\underline{mn}}\underset{2}{\mathcal{I}}_{ij\alpha}.\end{aligned}$$

Because of $P^{i}_{jk} = \overline{\Gamma}^{i}_{jk} - \Gamma^{i}_{jk}$ and $\underset{2}{\overline{\mathcal{I}}}_{ijk} = \underset{2}{\mathcal{I}}_{ijk}$ a following equation is a valid one:

$$\begin{aligned}&\underset{2}{\overline{\mathcal{I}}}_{ijm\bar{;}n} + \overline{\Gamma}^{\alpha}_{\underline{in}}\underset{2}{\overline{\mathcal{I}}}_{\alpha jm} + \overline{\Gamma}^{\alpha}_{\underline{jn}}\underset{2}{\overline{\mathcal{I}}}_{i\alpha n} + \overline{\Gamma}^{\alpha}_{\underline{mn}}\underset{2}{\overline{\mathcal{I}}}_{ij\alpha} \\ &\qquad = \underset{2}{\mathcal{I}}_{ijm;n} + \Gamma^{\alpha}_{\underline{in}}\underset{2}{\mathcal{I}}_{\alpha jm} + \Gamma^{\alpha}_{\underline{jn}}\underset{2}{\mathcal{I}}_{i\alpha n} + \Gamma^{\alpha}_{\underline{mn}}\underset{\mathcal{I}_{ij\alpha}}{}.\end{aligned} \tag{3.22}$$

Using the invariants $\underset{1}{\mathcal{I}}^{i}_{jk}, \underset{2}{\mathcal{I}}_{ijk}$ and the fact a composition of objects invariant of a mapping is an invariant of this mapping together with the equation (**??**), one concludes it is satisfied expressions:

$$\begin{aligned}
\overline{\Gamma}_{\underset{\vee}{i.jm;n}} &= \Gamma_{\underset{\vee}{i.jm;n}} - \underset{(1)}{\overline{\Gamma}}_{ijmn} + \underset{(1)}{\Gamma}_{ijmn},\\
\overline{\Gamma}^{\alpha}_{\underset{\vee}{jm}}\overline{\Gamma}_{\underset{\vee}{i.\alpha n}} &= \Gamma^{\alpha}_{\underset{\vee}{jm}}\Gamma_{\underset{\vee}{i.\alpha n}} - \underset{(2)}{\overline{\Gamma}}_{ijmn} + \underset{(2)}{\Gamma}_{ijmn},
\end{aligned} \tag{3.23}$$

for

$$\begin{aligned}
\underset{(1)}{\Gamma}_{ijmn} &= -\Big(\big(1-g^{-\frac{1}{N}}\big)\Gamma_{\underset{\vee}{i.jm}}\Big)_{;n} + \Big(\frac{1}{Ng^{\frac{1}{N}}}\big(g_{\underset{\vee}{ij}}\Gamma^{\alpha}_{\underline{m\alpha}} - g_{\underset{\vee}{im}}\Gamma^{\alpha}_{\underline{j\alpha}} + g_{\underset{\vee}{jm}}\Gamma^{\alpha}_{\underline{i\alpha}}\big)\Big)_{;n}\\
&\quad + \Gamma^{\alpha}_{\underline{in}}\underset{2}{\mathcal{I}}_{\alpha jm} + \Gamma^{\alpha}_{\underline{jn}}\underset{2}{\mathcal{I}}_{i\alpha n} + \Gamma^{\alpha}_{\underline{mn}}\underset{2}{\mathcal{I}}_{ij\alpha},\\
\underset{(2)}{\Gamma}_{ijmn} &= \big(1-g^{-\frac{1}{N}}\big)\Big(\Gamma^{\alpha}_{\underset{\vee}{jm}}\Gamma_{\underset{\vee}{\alpha.in}} - \frac{1}{2}\big(\Gamma^{\alpha}_{\underset{\vee}{\beta m}}A^{\beta}_{j} - \Gamma^{\alpha}_{\underset{\vee}{\beta j}}A^{\beta}_{m} - g^{\underline{\alpha\beta}}g_{\underset{\vee}{jm;\beta}}\big)\Big)\\
&\quad + \frac{1}{2}\Gamma_{\underset{\vee}{\alpha.in}}\big(\Gamma^{\alpha}_{\underset{\vee}{\beta m}}A^{\beta}_{j} - \Gamma^{\alpha}_{\underset{\vee}{\beta j}}A^{\beta}_{m} - g^{\underline{\alpha\beta}}g_{\underset{\vee}{jm;\beta}}\big)\\
&\quad + \frac{1}{Ng^{\frac{1}{N}}}\Gamma^{\alpha}_{\underset{\vee}{jm}}\big(g_{\underset{\vee}{i\alpha}}\Gamma^{\beta}_{\underline{n\beta}} - g_{\underset{\vee}{in}}\Gamma^{\beta}_{\underline{\alpha\beta}} + g_{\underset{\vee}{\alpha n}}\Gamma^{\beta}_{\underline{i\beta}}\big)\\
&\quad - \frac{1}{2Ng^{\frac{1}{N}}}\big(\Gamma^{\alpha}_{\underset{\vee}{\beta m}}A^{\beta}_{j} - \Gamma^{\alpha}_{\underset{\vee}{\beta j}}A^{\beta}_{m} - g^{\underline{\alpha\beta}}g_{\underset{\vee}{jm;\beta}}\big)\big(g_{\underset{\vee}{i\alpha}}\Gamma^{\gamma}_{\underline{n\gamma}} - g_{\underset{\vee}{in}}\Gamma^{\gamma}_{\underline{\alpha\gamma}} + g_{\underset{\vee}{\alpha n}}\Gamma^{\gamma}_{\underline{i\gamma}}\big).
\end{aligned}$$

Based on $\underset{k}{R}_{ij} = \underset{k}{R}^{\alpha}_{ij\alpha}$ and $\underset{k}{R} = g^{\underline{\alpha\beta}}\underset{k}{R}_{\alpha\beta}, k = 1,\ldots,5$, for curvature tensors (**??**-**??**), it is proved a following theorem is a valid one in this way:

Theorem 3.1 *Let $f : \mathbb{GR}_N \to \mathbb{G}\overline{\mathbb{R}}_N$ be a conformal mapping of a generalized Riemannian space $\mathbb{GR}_N$. Magnitudes:*

$$
\begin{aligned}
\underset{1}{C}_{ijmn} &= \underset{1}{R}_{ijmn} - \frac{1}{N-2}\big(g_{\underline{jm}}\underset{1}{R}_{in} - g_{\underline{jn}}\underset{1}{R}_{im} + g_{\underline{in}}\underset{1}{R}_{jm} - g_{\underline{im}}\underset{1}{R}_{jn}\big) \\
&\quad - \frac{\underset{1}{R}}{(N-1)(N-2)}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big) \\
&\quad - \frac{1}{N-2}\delta^{\epsilon}_{[i}\delta^{\rho}_{j]}\delta^{\alpha}_{[m}g_{\underline{\rho n}]}\big(\Gamma^{\beta}_{\underset{\vee}{\epsilon\alpha};\beta} + \Gamma^{\beta}_{\underset{\vee}{\epsilon\gamma}}\Gamma^{\gamma}_{\underset{\vee}{\alpha\beta}}\big) \\
&\quad + \frac{1}{(N-1)(N-2)}g^{\underline{\alpha\beta}}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big)\Gamma^{\gamma}_{\underset{\vee}{\alpha\delta}}\Gamma^{\delta}_{\underset{\vee}{\beta\gamma}} \\
&\quad + \underset{(1)}{\Gamma}_{ijmn} - \underset{(1)}{\Gamma}_{ijnm} + \underset{(2)}{\Gamma}_{ijmn} - \underset{(2)}{\Gamma}_{ijnm},
\end{aligned}
\tag{3.24}
$$

$$
\begin{aligned}
\underset{2}{C}_{ijmn} &= \underset{2}{R}_{ijmn} - \frac{1}{N-2}\big(g_{\underline{jm}}\underset{2}{R}_{in} - g_{\underline{jn}}\underset{2}{R}_{im} + g_{\underline{in}}\underset{2}{R}_{jm} - g_{\underline{im}}\underset{2}{R}_{jn}\big) \\
&\quad - \frac{\underset{2}{R}}{(N-1)(N-2)}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big) \\
&\quad + \frac{1}{N-2}\delta^{\epsilon}_{[i}\delta^{\rho}_{j]}\delta^{\alpha}_{[m}g_{\underline{\rho n}]}\big(\Gamma^{\beta}_{\underset{\vee}{\epsilon\alpha};\beta} - \Gamma^{\beta}_{\underset{\vee}{\epsilon\gamma}}\Gamma^{\gamma}_{\underset{\vee}{\alpha\beta}}\big) \\
&\quad + \frac{1}{(N-1)(N-2)}g^{\underline{\alpha\beta}}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big)\Gamma^{\gamma}_{\underset{\vee}{\alpha\delta}}\Gamma^{\delta}_{\underset{\vee}{\beta\gamma}} \\
&\quad - \underset{(1)}{\Gamma}_{ijmn} + \underset{(1)}{\Gamma}_{ijnm} + \underset{(2)}{\Gamma}_{ijmn} - \underset{(2)}{\Gamma}_{ijnm},
\end{aligned}
\tag{3.25}
$$

$$
\begin{aligned}
\underset{3}{C}_{ijmn} &= \underset{3}{R}_{ijmn} - \frac{1}{N-2}\big(g_{\underline{jm}}\underset{3}{R}_{in} - g_{\underline{jn}}\underset{3}{R}_{im} + g_{\underline{in}}\underset{3}{R}_{jm} - g_{\underline{im}}\underset{3}{R}_{jn}\big) \\
&\quad - \frac{\underset{3}{R}}{(N-1)(N-2)}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big) \\
&\quad - \frac{1}{N-2}\delta^{\epsilon}_{[i}\delta^{\rho}_{j]}\delta^{\alpha}_{[m}g_{\underline{\rho n}]}\big(\Gamma^{\beta}_{\underset{\vee}{\epsilon\alpha};\beta} + \Gamma^{\beta}_{\underset{\vee}{\epsilon\gamma}}\Gamma^{\gamma}_{\underset{\vee}{\alpha\beta}}\big) \\
&\quad + \frac{1}{(N-1)(N-2)}g^{\underline{\alpha\beta}}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big)\Gamma^{\gamma}_{\underset{\vee}{\alpha\delta}}\Gamma^{\delta}_{\underset{\vee}{\beta\gamma}} \\
&\quad + \underset{(1)}{\Gamma}_{ijmn} + \underset{(1)}{\Gamma}_{ijnm} - \underset{(2)}{\Gamma}_{ijmn} + \underset{(2)}{\Gamma}_{ijnm} - 2\underset{(2)}{\Gamma}_{imnj},
\end{aligned}
\tag{3.26}
$$

$$
\begin{aligned}
\underset{4}{C}_{ijmn} &= \underset{4}{R}_{ijmn} - \frac{1}{N-2}\big(g_{\underline{jm}}\underset{4}{R}_{in} - g_{\underline{jn}}\underset{4}{R}_{im} + g_{\underline{in}}\underset{4}{R}_{jm} - g_{\underline{im}}\underset{4}{R}_{jn}\big) \\
&\quad - \frac{\underset{4}{R}}{(N-1)(N-2)}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big) \\
&\quad - \frac{1}{N-2}\delta^{\epsilon}_{[i}\delta^{\rho}_{j]}\delta^{\alpha}_{[m}g_{\underline{\rho n}]}\big(\Gamma^{\beta}_{\underset{\vee}{\epsilon\alpha};\beta} - 3\Gamma^{\beta}_{\underset{\vee}{\epsilon\gamma}}\Gamma^{\gamma}_{\underset{\vee}{\alpha\beta}}\big) \\
&\quad + \frac{1}{(N-1)(N-2)}g^{\underline{\alpha\beta}}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big)\Gamma^{\gamma}_{\underset{\vee}{\alpha\delta}}\Gamma^{\delta}_{\underset{\vee}{\beta\gamma}} \\
&\quad + \underset{(1)}{\Gamma}_{ijmn} + \underset{(1)}{\Gamma}_{ijnm} - \underset{(2)}{\Gamma}_{ijmn} + \underset{(2)}{\Gamma}_{ijnm} + 2\underset{(2)}{\Gamma}_{imnj},
\end{aligned}
\tag{3.27}
$$

$$
\begin{aligned}
\underset{5}{C}_{ijmn} &= \underset{5}{R}_{ijmn} - \frac{1}{N-2}\big(g_{\underline{jm}}\underset{5}{R}_{in} - g_{\underline{jn}}\underset{5}{R}_{im} + g_{\underline{in}}\underset{5}{R}_{jm} - g_{\underline{im}}\underset{5}{R}_{jn}\big) \\
&\quad - \frac{\underset{5}{R}}{(N-1)(N-2)}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big) \\
&\quad - \frac{1}{N-2}\delta^{\epsilon}_{[i}\delta^{\rho}_{j]}\delta^{\alpha}_{[m}g_{\underline{\rho n}]}\Gamma^{\beta}_{\underset{\vee}{\epsilon\gamma}}\Gamma^{\gamma}_{\underset{\vee}{\alpha\beta}} \\
&\quad + \frac{1}{(N-1)(N-2)}g^{\underline{\alpha\beta}}\big(g_{\underline{im}}g_{\underline{jn}} - g_{\underline{in}}g_{\underline{jm}}\big)\Gamma^{\gamma}_{\underset{\vee}{\alpha\delta}}\Gamma^{\delta}_{\underset{\vee}{\beta\gamma}} \\
&\quad - \underset{(2)}{\Gamma}_{ijmn} + \underset{(2)}{\Gamma}_{ijnm},
\end{aligned}
\tag{3.28}
$$

for magnitudes $\underset{(1)}{\Gamma}_{ijmn}$ *and* $\underset{(2)}{\Gamma}_{ijmn}$ *defined above this theorem are invariants of the mapping* f. □

The invariants $\underset{\theta}{C}_{ijmn}, \theta = 1, \ldots, 5$, are **the θ-th generalized Weyl conformal curvature tensors**.

References

[1] **H. M. Abood**, *Almost Hermitian Manifold with Flat Bochner Tensor*, European Journal of Pure and Applied Mathematics, Vol. 3, No. 4, 2010, 730-736.

[2] **S. Bochner**, *Curvature and Betti Numbers. II*, 1949, Ann. of Math., Vol. 50, No. 1, pp. 77-93.

[3] **H. Chuda, M. Shiha**, *Conformal Holomorphically Projective Mappings Satisfying a Certain Initial Condition*, Miskolc Mathematical Notes, Vol. 14 (2013), No. 2, pp. 569-574.

[4] **L. P. Eisenhart**, *Riemannian Geometry*, Princeton University Press, 1997.

[5] **E. Goulart, M. Novello**, *Cosmological Stability of Weyl Conformal Tensor*, Gravitation and Cosmology, 2008, Vol. 14, No. 4, pp. 321-326.

[6] **S. Ivanov, M. Zlatanović**, *Connections on Non-symmetric (Generalized) Riemannian Manifold and Gravity*, arXiv, availible in: http://arxiv.org/pdf/1503.05217.pdf.

[7] **S. B. Mathur**, *Conformal spaces and curvature tensors*, Indian J. of Pure and Appl. Math., 2, No. 1, 1971, 29-31.

[8] **S. M. Minčić**, *Generalized Riemannian spaces*, Doctoral Thesis, Faculty of Sciences, University of Novi Sad, Novi Sad, (1975), (in Serbian).

[9] **S. M. Minčić**, *Ricci identities in the space of non-symmetric affine connection*, Matematički Vesinik, 10(25), Vol. 2, (1973), 161-172.

[10] **S. M. Minčić**, *Independent curvature tensors and pseudotensors of spaces with non-symmetric affine connexion*, Colloquia Mathematica Societatis Janos Bolayai, 31, Differential Geometry, Budapest (Hungary), (1979), 445-460.

[11] **P. Mocanu**, *Spatii partial projective*, Acad RPR, 6, 1955, 3, 4.

[12] **M. Prvanović**, *Projective and conformal transformations in recurrent and Ricci-recurent Riemannian space*, Tensor (NS), 12, No 3, 1962, 219-226.

[13] **N. S. Sinyukov**, *Geodesic Mappings of Riemannian Spaces*, "Nauka", Moscow, 1979, (in Russian).

[14] **M. S. Stanković, Lj. S. Velimirović, S. M. Minčić, M. Lj. Zlatanović**, *Equitorsion Conform Mappings of Generalized Riemannian Spaces*, Matematički Vesnik, 61 (2009), 119-129.

The author:
Nenad O. Vesić,
Faculty of Science and Mathematics address: Visegradska 33, 18000 Niš, Serbia